arXiv:1208.5310v1 [nlin.SI] 27 Aug 2012

Characteristic Lie rings and symmetries of differential Painlevé I and Painlevé III equations

O. S. Kostrigina[1]
Ufa State Aviation Technical University,
12, K. Marx str., Ufa, Russia, 450000
A. V. Zhiber[2]
Institute of mathematics with computer centre RAS,
112, Chernyshevsky str., Ufa, Russia, 450008

Abstract. *Two-component hyperbolic system of equations generated by ordinary differential Painlevé I*

$$u_{yy} = 6u^2 + y$$

and Painlevé III

$$yuu_{yy} = yu_y^2 - uu_y + \delta y + \beta u + \alpha u^3 + \gamma y u^4$$

equations are considered, where $\alpha, \beta, \gamma, \delta$ *are complex numbers. The structure of characteristic Lie rings is studied and higher symmetries of Lie-Bäcklund are obtained.*

Keywords: *characteristic vector field, characteristic ring,* $x-$ *and* $y-$*integrals, higher symmetries.*

Introduction

Characteristic Lie rings and higher symmetries of Lie-Bäcklund for Painlevé I

$$u_{yy} = 6u^2 + y \tag{1}$$

and Painlevé III

$$yuu_{yy} = yu_y^2 - uu_y + \delta y + \beta u + \alpha u^3 + \gamma y u^4 \tag{2}$$

equations are considered, where $\alpha, \beta, \gamma, \delta$ are complex numbers.

For definition of characteristic Lie ring of ordinary differential equations consider hyperbolic system of eqautions (see [1])

$$u^i_{xy} = F^i(x, y, u, u_x, u_y), \quad i = 1, 2, \ldots, n, \ u = (u^1, u^2, \ldots, u^n). \tag{3}$$

We introduce a set of independent variables $u_1 = u_x,\ \bar{u}_1 = u_y,\ u_2 = u_{xx},\ \bar{u}_2 = u_{yy}, \ldots$ and denote by $\bar{D}(D)$ an operator of full differentiation with respect to y (x).

---

[1] e-mail: kostrigina@mail.ru

[2] e-mail: zhiber@mail.ru

Definition 1 *Function* $\omega = \omega(x, y, u, u_1, \ldots, u_m)$ *is called an* $x-$*integral of order* m *of equations* (3) *if* $\sum_{i=1}^{n} \left(\frac{\partial \omega}{\partial u_m^i}\right)^2 \neq 0$ *and* $\bar{D}\omega = 0$. *Similarly,* $\bar{\omega} = \bar{\omega}(x, y, u, \bar{u}_1, \ldots, \bar{u}_p)$ *is an* $y-$*integral of order* p *of equations* (3), *if* $\sum_{i=1}^{n} \left(\frac{\partial \bar{\omega}}{\partial \bar{u}_p^i}\right)^2 \neq 0$ *and* $D\bar{\omega} = 0$.

Denote an area of locally analytical functions by $\Im$, where every function depends on a finite number of variables $x, y, \bar{u}_1, u, u_1, u_2, \ldots, u_k, \ldots$. Operator $\bar{D}$ acts on functions from $\Im$ as follows:

$$\bar{D} = \bar{u}_2^i X_i + X_{n+1},$$

where

$$X_i = \frac{\partial}{\partial \bar{u}_1^i}, \ i = 1, 2, \ldots, n,$$

$$X_{n+1} = \frac{\partial}{\partial y} + \bar{u}_1^i \frac{\partial}{\partial u^i} + F^i \frac{\partial}{\partial u_1^i} + D(F^i) \frac{\partial}{\partial u_2^i} + \ldots + D^{k-1}(F^i) \frac{\partial}{\partial u_k^i} + \ldots.$$

The $x-$characteristic Lie ring of equations (3) is a ring A generated by vector fields $X_1, X_2, \ldots, X_{n+1}$. Similarly, the $y-$characteristic Lie ring $\bar{A}$ is defined.

Notice that the concept of characteristic vector field first was introduced by the Goursat in [2].

Denote a linear area of commutator of length n, $n = 1, 2, 3, \ldots$ by L_n. For example L_1 is a linear span of vector fields $X_1, X_2, \ldots, X_{n+1}$ and L_2 is generated by operators $X_{ij} = [X_i, X_j], \ i, j = 1, 2, \ldots, n + 1$ etc. Then the $x-$ring is represented in the form

$$A = \sum_{i=1}^{\infty} L_i.$$

For $y-$characteristic ring $\bar{A}$ we have

$$\bar{A} = \sum_{i=1}^{\infty} \bar{L}_i.$$

In [3] it was conjectured that dimension of linear areas L_i ($\bar{L}_i$) for integrable equations grows slowly. Further this hypothesis was confirmed by numerous examples of integrable continuous and discrete models. Property of minimal growth became a kind of classification criterion for integrable equations. From [4] it is follows that this property of the ring and the property of existence of hierarchy of higher symmetries for integrable equations are equally universal.

Definition 2 *Symmetry of equations (3) is a set of functions*

$$f^i = f^i(x, y, u, u_1, \ldots, u_r, \bar{u}_1, \ldots, \bar{u}_s), \ i = 1, 2, \ldots, n,$$

satisfying the defining equations

$$D\bar{D}f^i = (F^i_{u^j} + F^i_{u_1^j} D + F^i_{\bar{u}_1^j}) f^j, \ i = 1, 2, \ldots, n.$$

It is known that any Lie-Bäcklund symmetry $f^i,\ i=1,2,\dots,n$ is represented in the form

$$f^i = f_1^i + f_2^i,$$

where functions f_1^i are depends on variables $x, y, u, u_1, u_2, \dots,$ and functions f_2^i are depends on variables $x, y, u, \bar{u}_1, \bar{u}_2, \dots$. And besides f_1^i and f_2^i are symmetries itself.

Concept of characteristic Lie ring for a system of ordinary differential equations

$$u_y^i = f^i(x, y, u), \quad i = 1, 2, \dots, n, \ u = (u^1, u^2, \dots, u^n), \tag{4}$$

was introduced in [5]. In this work two definitions for characteristic ring were proposed (4). First definition is based on change

$$u^i = \frac{\partial p^i}{\partial x}, \ i = 1, 2, \dots, n,$$

in which equations (4) are take the form

$$p_{xy}^i = f^i(x, y, v_x), \quad i = 1, 2, \dots, n, \ v = (v^1, v^2, \dots, v^n). \tag{5}$$

The x and $y-$characteristic Lie rings of hyperbolic system (5) are called characteristic Lie rings of initial system of ordinary differential equations (4).

Another definition of characteristic Lie ring for system (4) is associated with the following hyperbolic system of equations

$$u_{xy}^i = f_x^i + f_{u^k}^i u_x^k, \ i = 1, 2, \dots, n. \tag{6}$$

In the first section characteristic Lie ring for system of equations (5) and (6) corresponding to Painlevé I equation are considered. We show that these Lie rings are rings of slow growth. Namely, for the system of equations (5) dimension of $x-$ring is equal to three, and for $y-$ring $dim\bar{L}_1 = 3,\ dim\bar{L}_n = n,\ n = 2, 3, 4;\ dim\bar{L}_5 \leq 5$. For the hyperbolic system (6), corresponding to the equation Painlevé I, it is shown that dimension of $y-$ring is four, and for $x-$ring the following formalus are true $dimL_1 = 3,\ dimL_k = 1,\ k = 2, 3, 4, 5, 6;\ \ dim\sum_{i=1}^{6} L_i = 8$.

In the second and third sections the higher symmetries of Lie-Bäcklund for system of equations (5) corresponding to equations (1) and (2) are obtained .

1 Lie rings for Painlevé I equation

We rewrite equation (1) as

$$u_y = v, \quad v_y = 6u^2 + y.$$

Then the corresponding hyperbolic system of equations (5) and (6) take the form

$$p_{xy} = q_x, \quad q_{xy} = 6p_x^2 + y \ (u = p_x, \quad v = q_x) \tag{7}$$

and

$$u_{xy} = v_x, \quad v_{xy} = 12uu_x. \tag{8}$$

In this section we study the characteristic Lie ring of systems of equations (7) and (8).

For the system (7) $x-$characteristic Lie ring is generated by vector fields

$$X_1=\frac{\partial}{\partial p},\quad X_2=\frac{\partial}{\partial q},$$
$$X_3=\frac{\partial}{\partial y}+q_1\frac{\partial}{\partial p_1}+(6p_1^2+y)\frac{\partial}{\partial q_1}+q_2\frac{\partial}{\partial p_2}+12p_1p_2\frac{\partial}{\partial q_2}+\ldots.$$

Since $[X_1,X_2]=[X_1,X_3]=[X_2,X_3]=0$, then dimension of $x-$ring is equal to three. Herewith $x-$integrals

$$\omega=\omega(y,p_1,q_1) \text{ and } w=w(y,p_1,q_1)$$

are defined from partial derivative equations

$$\left(\frac{\partial}{\partial y}+q_1\frac{\partial}{\partial p_1}+(6p_1^2+y)\frac{\partial}{\partial q_1}\right)F=0. \tag{9}$$

Note, that $\omega=const$ and $w=const$ are define integrals of initial equation (1).

The $y-$characteristic Lie ring of equations (7) is defined by vector fields

$$\begin{gathered}
Y_1=\frac{\partial}{\partial p_1},\quad Y_2=\frac{\partial}{\partial q_1},\\
Y_3=\frac{\partial}{\partial x}+p_1\frac{\partial}{\partial p}+q_1\frac{\partial}{\partial \lambda_1}+(6p_1^2+y)\frac{\partial}{\partial \lambda_2}+(12p_1q_1+1)\frac{\partial}{\partial \lambda_3}+\\
+(12q_1^2+72p_1^3+12yp_1)\frac{\partial}{\partial \lambda_4}+(360q_1p_1^2+12p_1+36yq_1)\frac{\partial}{\partial \lambda_5}+\\
+(2160p_1^4+720p_1q_1^2+48q_1+576yp_1^2+36y^2)\frac{\partial}{\partial \lambda_6}+\ldots,
\end{gathered} \tag{10}$$

where

$$\frac{\partial}{\partial \lambda_i}=\frac{\partial}{\partial \bar{p}_i}+\frac{\partial}{\partial \bar{q}_{i-1}},\ i=1,2,\ldots.$$

It is easy to see, that the vector fields $Y_1,\ [Y_1,Y_3],\ [Y_1,[Y_1,Y_3]],\ [Y_1,[Y_1,[Y_1,Y_3]]],\ldots$ are linearly independent, and hence $y-$ring is infinite.

Since D and $\bar{D}$ commute, we have

$$\begin{gathered}
[\bar{D},D]F(x,y,p,q,p_1,q_1,\bar{p}_1,\bar{q}_1,\bar{p}_2,\bar{q}_2,\ldots)=\\
=\bar{D}(p_2Y_1+q_2Y_2+Y_3)F-(p_2Y_1+q_2Y_2+Y_3)\bar{D}F=\\
=(p_2[Y,Y_1]+q_2[Y,Y_2]+[Y,Y_3]+q_2Y_1+12p_1p_2Y_2)F=0.
\end{gathered} \tag{11}$$

Here Y is the operator of full differentiation with respect to y in the area of functions depending on a finite set of variables $x,y,p,q,p_1,q_1,\bar{p}_1,\bar{q}_1,\bar{p}_2,\bar{q}_2,\ldots$

$$Y=\frac{\partial}{\partial y}+\bar{p}_1\frac{\partial}{\partial p}+\bar{q}_1\frac{\partial}{\partial q}+q_1\frac{\partial}{\partial p_1}+(6p_1^2+y)\frac{\partial}{\partial q_1}+\bar{p}_2\frac{\partial}{\partial \bar{p}_1}+\bar{q}_2\frac{\partial}{\partial \bar{q}_1}+\bar{p}_3\frac{\partial}{\partial \bar{p}_2}+\bar{q}_3\frac{\partial}{\partial \bar{q}_2}+\ldots.$$

The following statement is true.

Lemma 1 *Let the vector field $\bar{Z}$ is given by*

$$\bar{Z} = \sum_{i=1}^{\infty} \left(\alpha_i \frac{\partial}{\partial \bar{p}_i} + \beta_i \frac{\partial}{\partial \bar{q}_i} \right),$$

$$\alpha_i = \alpha_i(y, p_1, q_1), \ \beta_i = \beta_i(y, p_1, q_1).$$

Then the equality $[Y, \bar{Z}] = 0$ is true if and only if $\bar{Z} = 0$.

Proof. We have

$$[Y, \bar{Z}] = \sum_{i=1}^{\infty} \left(Y(\alpha_i) \frac{\partial}{\partial \bar{p}_i} + Y(\beta_i) \frac{\partial}{\partial \bar{q}_i} \right) - \sum_{i=1}^{\infty} \left(\alpha_i \frac{\partial}{\partial \bar{p}_{i-1}} + \beta_i \frac{\partial}{\partial \bar{q}_{i-1}} \right) = 0,$$

that is $\alpha_1 = 0, \ \beta_1 = 0, \ \alpha_{i+1} = Y(\alpha_i), \ \beta_{i+1} = Y(\beta_i), \ i = 1, 2, \ldots$. Hence $\alpha_i = 0$, $\beta_i = 0, i = 1, 2, \ldots$ and $\bar{Z} = 0$.

Lemma 1 is proved.

From (11) it is follows that

$$[Y, Y_1] = -12 p_1 Y_2, \ [Y, Y_2] = -Y_1, \ [Y, Y_3] = 0. \tag{12}$$

The linear area $\bar{L}_2$ is generated by operators Y_{13} and Y_{23} :

$$Y_{13} = \frac{\partial}{\partial p} + 12 p_1 \frac{\partial}{\partial \lambda_2} + 12 q_1 \frac{\partial}{\partial \lambda_3} + (216 p_1^2 + 12 y) \frac{\partial}{\partial \lambda_4} + (720 p_1 q_1 + 12) \frac{\partial}{\partial \lambda_5} +$$
$$+ (8640 p_1^3 + 720 q_1^2 + 1152 y p_1) \frac{\partial}{\partial \lambda_6} + \ldots, \tag{13}$$
$$Y_{23} = \frac{\partial}{\partial \lambda_1} + 12 p_1 \frac{\partial}{\partial \lambda_3} + 24 q_1 \frac{\partial}{\partial \lambda_4} + (360 p_1^2 + 36 y) \frac{\partial}{\partial \lambda_5} + (1440 p_1 q_1 + 48) \frac{\partial}{\partial \lambda_6} + \ldots,$$

Using the Jacobi identity, we obtain

$$[Y, Y_{13}] = -[Y_3, [Y, Y_1]] + [Y_1, [Y, Y_3]], \quad [Y, Y_{23}] = -[Y_3, [Y, Y_2]] + [Y_2, [Y, Y_3]],$$

or, using (12), we have

$$[Y, Y_{13}] = -[Y_3, -12 p_1 Y_2], \quad [Y, Y_{23}] = -[Y_3, -Y_1].$$

Thus, we have the following relations

$$[Y, Y_{13}] = -12 p_1 Y_{23}, \quad [Y, Y_{23}] = -Y_{13}. \tag{14}$$

The linear area $\bar{L}_3$ is generated by commutators $Y_{113}, \ Y_{123}, \ Y_{213}, \ Y_{223}$. Herewith

$$Y_{213} = Y_{123}$$

and

$$Y_{113} = 12 \frac{\partial}{\partial \lambda_2} + 432 p_1 \frac{\partial}{\partial \lambda_4} + 720 q_1 \frac{\partial}{\partial \lambda_5} + (25920 p_1^2 + 1152 y) \frac{\partial}{\partial \lambda_6} + \ldots,$$
$$Y_{123} = 12 \frac{\partial}{\partial \lambda_3} + 720 p_1 \frac{\partial}{\partial \lambda_5} + 1440 q_1 \frac{\partial}{\partial \lambda_6} + \ldots, \tag{15}$$
$$Y_{223} = 24 \frac{\partial}{\partial \lambda_4} + 1440 p_1 \frac{\partial}{\partial \lambda_6} + \ldots.$$

As above, we can prove the following formulas

$$[Y,Y_{213}]=-Y_{113}-12p_1Y_{223},\quad [Y,Y_{113}]=-12Y_{23}-24p_1Y_{123},\quad [Y,Y_{223}]=-2Y_{123}. \tag{16}$$

The linear area $\bar{L}_4$ is generated by operators $Y_{1113},\ Y_{2113},\ Y_{1213},\ Y_{2213},\ Y_{1223},\ Y_{2223}$, $[Y_{13},Y_{23}]$, herewith

$$Y_{1213}=Y_{2113},\ Y_{1223}=Y_{2213},\ [Y_{13},Y_{23}]=0$$

and

$$\begin{gathered}Y_{1113}=432\frac{\partial}{\partial\lambda_4}+51840p_1\frac{\partial}{\partial\lambda_6}+\ldots,\\ Y_{2113}=720\frac{\partial}{\partial\lambda_5}+0\cdot\frac{\partial}{\partial\lambda_6}+\ldots,\\ Y_{2213}=1440\frac{\partial}{\partial\lambda_6}\ldots,\quad Y_{2223}=0\cdot\frac{\partial}{\partial\lambda_6}+\ldots.\end{gathered} \tag{17}$$

We can show that

$$[Y,Y_{1113}]=-36p_1Y_{2113}-36Y_{123},\quad [Y,Y_{2213}]=-2Y_{2113}-12p_1Y_{2223}. \tag{18}$$

From (10), (13), (15) it is follows that $dim\bar{L}_1=3,\ dim\bar{L}_2=2,\ dim\bar{L}_3=3$. Let us show that the operators $Y_{223},Y_{1113},Y_{2113},Y_{2213},Y_{2223}$ are linearly independent. If they are dependent then according to (15), (17) we must have

$$Y_{1113}-\frac{432}{24}Y_{223}=\frac{25920}{1440}p_1Y_{2213}.$$

According to Lemma 1, this relation is equivalent to

$$[Y,Y_{1113}]-18[Y,Y_{223}]=18q_1Y_{2213}+18p_1[Y,Y_{2213}].$$

Using the formulas (16), (18), we have

$$-36p_1Y_{2113}-36Y_{123}+36Y_{123}=18q_1Y_{2213}+18p_1(-2Y_{2113}-12p_1Y_{2223})$$

or

$$q_1Y_{2213}-12p_1^2Y_{2223}=0.$$

That it is impossible. Hence operators $Y_{223},Y_{1113},Y_{2113},Y_{2213},Y_{2223}$ are linearly independent and $dim\bar{L}_4=4$.

The area $\bar{L}_5$ is generated by commutators $Y_{11113},\ Y_{21113},\ Y_{12113},\ Y_{22113},\ Y_{12213},\ Y_{22213}$, $Y_{12223},\ Y_{22223}$, and among them there are pairs of equal operators. Namely $Y_{21113}=Y_{12113},\ Y_{22113}=Y_{12213},\ Y_{22213}=Y_{12223}$. Thus $dim\bar{L}_5\leq 5$.

Based on the above results, it seems possible to assume the validity of the following formulas

$$dim\bar{L}_k\leq k,\ k\geq 6.$$

Then we have

$$dim(\bar{L}_1+\bar{L}_2+\ldots+\bar{L}_n)\leq 3+2+3+4+5+\sum_{k=6}^{n}k,$$

or

$$dim(\bar{L}_1+\bar{L}_2+\ldots+\bar{L}_n)\leq\frac{n^2}{2}+\frac{n}{2}+2.$$

Thus, we can hypothesize that the $y-$ring of (7) is a ring of slow growth.

Next, we consider the characteristic Lie ring of equations (8).

$Y-$characteristic Lie ring of (8) is generated by operators

$$Y_1=\frac{\partial}{\partial u_1},\ Y_2=\frac{\partial}{\partial v_1},$$

$$Y_3=u_1\left(\frac{\partial}{\partial u}+12u\frac{\partial}{\partial\bar{v}_1}+12u\frac{\partial}{\partial\bar{u}_2}+12\bar{u}_1\frac{\partial}{\partial\bar{v}_2}+\ldots\right)+v_1\left(\frac{\partial}{\partial v}+\frac{\partial}{\partial\bar{u}_1}+12u\frac{\partial}{\partial\bar{v}_2}+\ldots\right).$$

Clearly that $Y_3=u_1Y_{13}+v_1Y_{23}$ and operators $Y_1,\ Y_2,\ Y_{13},\ Y_{23}$ form a basis of $y-$characteristic ring. Thus, the dimension of $y-$ring is equal to 4. And equations (8) has two $y-$integrals of the first order:

$$\bar{\omega}=\bar{u}_1-v,\ \bar{w}=\bar{v}_1-6u^2.$$

Consider $x-$characteristic ring. The operator of full differentiation with respect to y in the area of functions depending on a set of variables $u,v,u_1,v_1,u_2,v_2,\ldots$ is given by

$$\bar{D}=\bar{u}_1X_1+\bar{v}_1X_2+X_3,$$

where

$$X_1=\frac{\partial}{\partial u},\ X_2=\frac{\partial}{\partial v},$$

$$X_3=\left(v_1\frac{\partial}{\partial u_1}+v_2\frac{\partial}{\partial u_2}+v_3\frac{\partial}{\partial u_3}+\ldots\right)+$$
$$+12\left(uu_1\frac{\partial}{\partial v_1}+(uu_2+u_1^2)\frac{\partial}{\partial v_2}+(uu_3+3u_1u_2)\frac{\partial}{\partial v_3}+\ldots\right).$$

Since operators D and $\bar{D}$ commute, we have

$$\begin{gathered}[\bar{D},D]F(u,v,u_1,v_1,u_2,v_2,\ldots)=\\=(D(\bar{u}_1X_1+\bar{v}_1X_2+X_3)-(\bar{u}_1X_1+\bar{v}_1X_2+X_3)D)F=\\=(\bar{u}_1[D,X_1]+\bar{v}_1[D,X_2]+[D,X_3]+v_1X_1+12uu_1X_2)F=0.\end{gathered}\tag{19}$$

Here X is the operator of full differentiation with respect to x in the area of functions depending on a set of variables $u,v,u_1,v_1,u_2,v_2,\ldots$

$$X=u_1\frac{\partial}{\partial u}+v_1\frac{\partial}{\partial v}+u_2\frac{\partial}{\partial u_1}+v_2\frac{\partial}{\partial v_1}+\ldots.$$

For the operator X we can prove the following statement similarly as in the lemma 1.

Lemma 2 *Let the vector field Z is given by*

$$Z = \sum_{i=1}^{\infty} \left(\delta_i \frac{\partial}{\partial u_i} + \epsilon_i \frac{\partial}{\partial v_i} \right),$$

$$\delta_i = \delta_i(u, v, u_1, v_1, u_2, v_2, \ldots, u_{n_i}, v_{n_i}), \;\; \epsilon_i = \epsilon_i(u, v, u_1, v_1, u_2, v_2, \ldots, u_{k_i}, v_{k_i}).$$

Then $[X, Z] = 0$ if and only if $Z = 0$.

From (19) it is follows

$$[X, X_1] = 0, \;\; [X, X_2] = 0, \;\; [X, X_3] = -v_1 X_1 - 12uu_1 X_2. \tag{20}$$

It is easy to see that the dimension of the linear area L_1 ($L_1 = L\langle X_1, X_2, X_3 \rangle$) is equal to three.

Since the coefficients of the vector field X_3 are independent of v, then commutator $X_{23} = 0$. And commutator X_{13} has the form

$$X_{13} = 12 \left(u_1 \frac{\partial}{\partial v_1} + u_2 \frac{\partial}{\partial v_2} + u_3 \frac{\partial}{\partial v_3} + u_4 \frac{\partial}{\partial v_4} + \ldots \right).$$

Thus $dim L_2 = 1$. Herewith

$$[X, X_{13}] = -[X_3, [X, X_1]] + [X_1, [X, X_3]],$$

or, using (20), we have

$$[X, X_{13}] = [X_1, -v_1 X_1 - 12uu_1 X_2],$$

that is

$$[X, X_{13}] = -12u_1 X_2. \tag{21}$$

The form of the operators X_1, X_2, X_3 and X_{13} implies that X_{113} and X_{213} are zero, and the commutator X_{313} has the form

$$X_{313} = 12 \left(v_1 \frac{\partial}{\partial v_1} + v_2 \frac{\partial}{\partial v_2} + v_3 \frac{\partial}{\partial v_3} + v_4 \frac{\partial}{\partial v_4} + \ldots \right) -$$
$$-12 \left(u_1 \frac{\partial}{\partial u_1} + u_2 \frac{\partial}{\partial u_2} + u_3 \frac{\partial}{\partial u_3} + u_4 \frac{\partial}{\partial u_4} + \ldots \right).$$

Taking into account (20), (21), we have

$$[X, X_{313}] = -[X_{13}, [X, X_3]] + [X_3, [X, X_{13}]] = -[X_{13}, -v_1 X_1 - 12uu_1 X_2] + [X_3, -12u_1 X_2],$$

or

$$[X, X_{313}] = 12u_1 X_1 - 12v_1 X_2. \tag{22}$$

Let show that the operators X_3, X_{13} and X_{313} are linearly independent. Indeed, if they are linearly dependent, then there exists functions α, β of the variables $u, v, u_1, v_1, u_2, v_2, \ldots$ such that $X_{313} = \alpha X_3 + \beta X_{13}$. Last equality, according to Lemma 2, is equivalent to

$$[X, X_{313}] = D(\alpha) X_3 + D(\beta) X_{13} + \alpha [X, X_3] + \beta [X, X_{13}].$$

Or using (20), (21), (22), we have

$$12u_1X_1 - 12v_1X_2 = X(\alpha)X_3 + X(\beta)X_{13} + \alpha(-v_1X_1 - 12uu_1X_2) + \beta(-12u_1X_2).$$

Equating the coefficients of the independent operators X_1 and X_3, we obtain the system of equations

$$X(\alpha) = 0,\ 12u_1 = -v_1\alpha.$$

The first equality implies that $\alpha = const$. This contradicts the second equation. Consequently, the operators $X_3,\ X_{13}$ and X_{313} are linearly independent and $dimL_3 = 1$.

The linear area L_4 is generated by the operators $X_{1313},\ X_{2313}$ and X_{3313}. Thus, it is easy to see that $X_{1313} = X_{2313} = 0$, and the commutator X_{3313} has the form

$$X_{3313} = -24v_1\frac{\partial}{\partial u_1} + 288uu_1\frac{\partial}{\partial v_1} + \ldots.$$

Using formulas (20), (22), we find

$$[X, X_{3313}] = -[X_{313}, [X, X_3]] + [X_3, [X, X_{313}]] =$$
$$= -[X_{313}, -v_1X_1 - 12uu_1X_2] + [X_3, 12u_1X_1 - 12v_1X_2],$$

hence

$$[X, X_{3313}] = 24v_1X_1 - 288uu_1X_2 - 12u_1X_{13}. \tag{23}$$

Let show that the operators $X_3,\ X_{13},\ X_{313}$ and X_{3313} are linearly independent. Suppose to the contrary, that $X_{3313} = \alpha X_3 + \beta X_{13} + \delta X_{313}$, where $\alpha,\ \beta,\ \delta$ is a function of the variables $u, v, u_1, v_1, u_2, v_2, \ldots$. According to Lemma 2 and formulas (20) - (23), we have

$$24v_1X_1 - 288uu_1X_2 - 12u_1X_{13} = X(\alpha)X_3 + X(\beta)X_{13} + X(\delta)X_{313} +$$
$$+\alpha(-v_1X_1 - 12uu_1X_2) + \beta(-12u_1X_2) + \delta(12u_1X_1 - 12v_1X_2).$$

Equating coefficients of the operators $X_3,\ X_{313},\ X_1$ and $X_2,\ X_{13}$ in the resulting correlation, we have

$$X(\alpha) = 0,\ X(\delta) = 0,\ \ 24v_1 = -v_1\alpha + 12u_1\delta,$$
$$-288uu_1 = -12uu_1\alpha - 12u_1\beta - 12v_1\delta,\ -12u_1 = X(\beta).$$

From the first three equations follows that $\alpha = -24,\ \delta = 0$. Substituting the values α and δ in the fourth equation we find that $\beta = 48u$. This contradicts the condition $-12u_1 = X(\beta)$. Hence $dimL_4 = 1$.

The area L_5 is the linear span of commutators $X_{13313},\ X_{23313},\ X_{33313},\ [X_{13}, X_{313}]$. Using the Jacobi identity and the formulas (21) - (23), we can show that

$$[X, X_{13313}] = -288u_1X_2,\ [X, X_{23313}] = 0,\ [X, [X_{13}, X_{313}]] = -288u_1X_2.$$

Then, according to Lemma 2 and equality (21), we obtain

$$X_{13313} = 24X_{13},\ X_{23313} = 0,\ [X_{13}, X_{313}] = 24X_{13}.$$

For operator X_{33313} we have

$$[X, X_{33313}] = -[X_{3313}, [X, X_3]] + [X_3, [X, X_{3313}]] =$$
$$= -[X_{3313}, -v_1X_1 - 12uu_1X_2] + [X_3, 24v_1X_1 - 288uu_1X_2 - 12u_1X_{13}].$$

Rearranging this equation, we obtain the equality

$$[X, X_{33313}] = 576uu_1X_1 - 576uv_1X_2 - 60v_1X_{13} - 12u_1X_{313}.$$

Using this equality we can obtain, that the operators $X_3,\ X_{13},\ X_{313},\ X_{3313}$ and X_{33313} are linearly independent, and hence $dimL_5 = 1$.

Finally, we consider the area $L_6 = L\left\langle X_{133313},\ X_{233313},\ X_{333313},\ [X_{13}, X_{3313}]\right\rangle$. Easy to check the validity of the following formulas

$$X_{233313} = 0,\ X_{133313} = 48X_{313},\ [X_{13}, X_{3313}] = 24X_{313},$$
$$[X, X_{333313}] = 1152uv_1X_1 + 6912u^2u_1X_2 - 1296uu_1X_{13} - 120v_1X_{313} - 12u_1X_{3313}.$$

As above, we can prove that the operators $X_3,\ X_{13},\ X_{313},\ X_{3313}$ and X_{333313} are linearly independent and $dimL_6 = 1$.

Thus, we have shown that

$$dimL_k = 1,\ k = 2, 3, 4, 5, 6;\quad dim\sum_{i=1}^{n} L_i = 8,\ n = 6.$$

Apparently, these formulas are valid for any k and n, ie

$$dimL_k = 1,\ k \geq 2;\quad dim\sum_{i=1}^{n} L_i = n + 2,$$

and $x-$ring of the system of equations (8) is a ring of slow growth.

2 Symmetries of Painlevé I equation

In this section we calculate the higher symmetry of the system (7).

Consider $x-$symmetries of the form

$$f = f(x, y, p, q, p_1, q_1, \ldots, p_n, q_n),\quad g = g(x, y, p, q, p_1, q_1, \ldots, p_n, q_n),$$
$$(p_\tau = f,\ q_\tau = g).$$

We move from variables $x, y, p, q, p_1, q_1, \ldots, p_n, q_n, \ldots$ to $x, y, p, q\ \omega,\ w,\ \omega_1,\ w_1, \ldots,\ \omega_{n-1}$, $w_{n-1}, \ldots$. Then the symmetries of f and g can be written as

$$f = f(x, y, p, q, \omega, w, \omega_1, w_1, \ldots, \omega_{n-1}, w_{n-1}),\ \ g = g(x, y, p, q, \omega, w, \omega_1, w_1, \ldots, \omega_{n-1}, w_{n-1}),$$

where ω and w are $x-$integrals of the first order of equations (7) and $\omega_k = D^k\omega,\ w_k = D^kw,\ k = 1, 2, \ldots$.

The defining system of equations has the following view

$$D\bar{D}f = Dg, \quad D\bar{D}g = 12p_1 Df. \tag{24}$$

Further we introduce the notation

$$\begin{aligned} Df &= F(x, y, p, q, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n), \\ Dg &= G(x, y, p, q, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n). \end{aligned} \tag{25}$$

Then equations (24) become

$$\bar{D}F = G, \quad \bar{D}G = 12p_1 F. \tag{26}$$

From (25), (26) we get that $F_p = F_q = G_p = G_q = 0$ and, hence equations (26) are equivalent to the following

$$F_y = G, \quad G_y = 12p_1 F. \tag{27}$$

Since equations (7) have $x-$integrals of the first order, then there exists a function $h = h(y, \omega, w)$ such that $p_1 = h$. Thus from (27) we get

$$F_{yy} = 12h(y, \omega, w)F. \tag{28}$$

Consider the symmetry of the first order

$$f = p_1, \ g = q_1.$$

We have

$$F = Df = Dh = h_\omega \omega_1 + h_w w_1,$$

and equation (28) becomes

$$h_{\omega yy}\omega_1 + h_{wyy}w_1 = 12h(h_\omega \omega_1 + h_w w_1).$$

Which implies that

$$h_{\omega yy} = 12hh_\omega, \quad h_{wyy} = 12hh_w.$$

Hence the functions h_ω and h_w are particular solutions of (28). We show that the solutions h_ω and h_w are linearly independent. Indeed, if $h_\omega = c(\omega, w)h_w$, then $h = h(y, \alpha(\omega, w)) = h(y, W)$. And, therefore, system (7) has an integral $W = W(y, p_1)$, which is impossible because of equality (9).

Thus, the general solution of (28) has the form

$$F = h_\omega A(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n) + h_w B(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n). \tag{29}$$

Finally from equations (25), (27), (29) it follows that the higher local $x-$Lie-Bücklund symmetries of (7) are given by

$$f = D^{-1}(h_\omega A + h_w B), \quad g = D^{-1}(h_{\omega y} A + h_{wy} B),$$

where functions $A = A(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n)$ and $B = B(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n)$ satisfy

$$\frac{\delta}{\delta\omega}(h_\omega A + h_w B) = \frac{\delta}{\delta w}(h_\omega A + h_w B) = 0.$$

Further we construct the $y-$symmetries

$$\varphi = \varphi(x, y, p, q, \bar{p}_1, \bar{q}_1, \ldots, \bar{p}_n, \bar{q}_n), \quad \psi = \psi(x, y, p, q, \bar{p}_1, \bar{q}_1, \ldots, \bar{p}_m, \bar{q}_m),$$
$$(p_\tau = \varphi, \ q_\tau = \psi)$$

for system (7).

Let the order of variables $p, q, \bar{p}_1, \bar{q}_1, \bar{p}_2, \bar{q}_2, \ldots$ the functions $D\varphi$ and $D\psi$ be equal to n and m correspondingly. Then, from the defining equations

$$D\bar{D}\varphi = D\psi, \quad D\bar{D}\psi = 12p_1 D\varphi \tag{30}$$

it follows that $n+1 = m$ и $m+1 = n$ and hence

$$D\varphi = F(x, y, p_1, q_1), \quad D\psi = G(x, y, p_1, q_1).$$

On the other hand

$$D\varphi(x, y, p, q, \bar{p}_1, \bar{q}_1, \ldots, \bar{p}_n, \bar{q}_n) = \frac{\partial\varphi}{\partial x} + p_1\frac{\partial\varphi}{\partial p} + q_1\left(\frac{\partial}{\partial\bar{p}_1} + \frac{\partial}{\partial q}\right)\varphi +$$
$$+(6p_1^2 + y)\left(\frac{\partial}{\partial\bar{p}_2} + \frac{\partial}{\partial\bar{q}_1}\right)\varphi + (12p_1q_1 + 1)\left(\frac{\partial}{\partial\bar{p}_3} + \frac{\partial}{\partial\bar{q}_2}\right)\varphi + \ldots.$$

Hence the function φ satisfies the equations

$$\frac{\partial\varphi}{\partial x} = \alpha(x, y), \ \frac{\partial\varphi}{\partial p} = \alpha_0(x, y), \ \left(\frac{\partial}{\partial\bar{p}_1} + \frac{\partial}{\partial q}\right)\varphi = \alpha_1(x, y),$$
$$\left(\frac{\partial}{\partial\bar{p}_2} + \frac{\partial}{\partial\bar{q}_1}\right)\varphi = \alpha_2(x, y), \ldots, \ \left(\frac{\partial}{\partial\bar{p}_n} + \frac{\partial}{\partial\bar{q}_{n-1}}\right)\varphi = \alpha_n(x, y).$$

It is easy to obtain from it, that

$$\varphi = \beta(x, y) + \beta_0(y)p + \beta_1(y)\bar{p}_1 + \ldots + \beta_n(y)\bar{p}_n + h(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{n-1}), \tag{31}$$

where $\bar{\omega} = \bar{p}_1 - q$ is the $y-$integral of system (7).
An analogous formula holds true for the function ψ :

$$\psi = \gamma(x, y) + \gamma_0(y)p + \gamma_1(y)\bar{p}_1 + \ldots + \gamma_m(y)\bar{p}_m + H(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{m-1}). \tag{32}$$

Since the functions $\varphi = h(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{n-1})$ and $\psi = H(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{m-1})$ are define symmetries of the system (7) for any h and H, then we obtain from (31) and (32), that

$$\varphi = \beta(x, y) + \sum_{k=0}^{n}\beta_k(y)\bar{p}_k, \quad \psi = \gamma(x, y) + \sum_{k=0}^{m}\gamma_k(y)\bar{p}_k \tag{33}$$

are also symmetries. Substituting (33) into defining system (30), we find, that

$$\varphi = \varphi(y), \quad \psi = \psi(y).$$

Thus $y-$symmetries of the equations (7) are calculated by formulae

$$\varphi = h(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{n-1}), \quad \psi = H(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{m-1}).$$

3 Symmetries of Painlevé III equation

System of equations (5) corresponding to Painlevé III equation have the form

$$p_{xy} = q_x, \quad y p_x q_{xy} = y q_x^2 - p_x q_x + \delta y + \beta p_x + \alpha p_x^3 + \gamma y p_x^4. \tag{34}$$

In this section higher symmetries for the system (34) are built.

The $x-$characteristic Lie ring of system (34) is generated by vector fields

$$X_1 = \frac{\partial}{\partial p}, \quad X_2 = \frac{\partial}{\partial q},$$
$$X_3 = \frac{\partial}{\partial y} + q_1 \frac{\partial}{\partial p_1} + \left(\frac{q_1^2}{p_1} - \frac{q_1}{y} + \delta \frac{1}{p_1} + \beta \frac{1}{y} + \alpha \frac{p_1^2}{y} + \gamma p_1^3 \right) \frac{\partial}{\partial q_1} + \dots .$$

Since the coefficients of the vector field X_3 is independent of p and q, the dimension of $x-$ring is equal to three and $x-$integrals $\omega = \omega(y, p_1, q_1)$ and $w = w(y, p_1, q_1)$ are defined by the partial differential equation

$$\left(\frac{\partial}{\partial y} + q_1 \frac{\partial}{\partial p_1} + \left(\frac{q_1^2}{p_1} - \frac{q_1}{y} + \delta \frac{1}{p_1} + \beta \frac{1}{y} + \alpha \frac{p_1^2}{y} + \gamma p_1^3 \right) \frac{\partial}{\partial q_1} \right) \Phi = 0.$$

Note, that $\omega = const$ and $w = const$ are define the integrals of the initial equation (2).

Consider higher $x-$ symmetries for equations (34)

$$f = f(x, y, p, q, p_1, q_1, \dots, p_n, q_n), \quad g = g(x, y, p, q, p_1, q_1, \dots, p_n, q_n),$$
$$(p_\tau = f, \; q_\tau = g).$$

We move from variables $x, y, p, q, p_1, q_1, \dots, p_n, q_n, \dots$ to x, y, p, q ω, w, ω_1, $w_1, \dots, \omega_{n-1}$, $w_{n-1}, \dots$. Then the symmetry of f and g can be written as

$$f = f(x, y, p, q, \omega, w, \omega_1, w_1, \dots, \omega_{n-1}, w_{n-1}), \; g = g(x, y, p, q, \omega, w, \omega_1, w_1, \dots, \omega_{n-1}, w_{n-1}).$$

The defining system for system (34) has the form

$$D\bar{D}f = Dg, \quad D\bar{D}g = \left(2\frac{q_1}{p_1} - \frac{1}{y} \right) Dg + \left(-\frac{q_1^2}{p_1^2} - \delta \frac{1}{p_1^2} + 2\alpha \frac{p_1}{y} + 3\gamma p_1^2 \right) Df, \tag{35}$$

which can be written as

$$\bar{D}F = G, \quad \bar{D}G = \left(2\frac{q_1}{p_1} - \frac{1}{y} \right) G + \left(-\frac{q_1^2}{p_1^2} - \delta \frac{1}{p_1^2} + 2\alpha \frac{p_1}{y} + 3\gamma p_1^2 \right) F, \tag{36}$$

where

$$\begin{aligned} Df &= F(x, y, p, q, \omega, w, \omega_1, w_1, \dots, \omega_n, w_n), \\ Dg &= G(x, y, p, q, \omega, w, \omega_1, w_1, \dots, \omega_n, w_n). \end{aligned} \tag{37}$$

Clearly that $F_p = F_q = G_p = G_q = 0$ and the system (36) is equivalent to

$$F_y = G, \quad G_y = \left(2\frac{q_1}{p_1} - \frac{1}{y} \right) G + \left(-\frac{q_1^2}{p_1^2} - \delta \frac{1}{p_1^2} + 2\alpha \frac{p_1}{y} + 3\gamma p_1^2 \right) F. \tag{38}$$

Since the system (34) has $x-$ integrals of the first order, then there are functions $h = h(y, \omega, w)$ and $H = H(y, \omega, w)$ such that $p_1 = h,\ q_1 = H$. Then the second equation (38) can be written as

$$F_{yy} = \left(2\frac{H}{h} - \frac{1}{y}\right) F_y + \left(-\frac{H^2}{h^2} - \delta\frac{1}{h^2} + 2\alpha\frac{h}{y} + 3\gamma h^2\right) F. \tag{39}$$

For the symmetry of the first order

$$f = p_1,\ g = q_1$$

equation (39) is transformed as follows

$$h_{\omega yy}\omega_1 + h_{wyy}w_1 = \left(2\frac{H}{h} - \frac{1}{y}\right)(h_{\omega y}\omega_1 + h_{wy}w_1) + \left(-\frac{H^2}{h^2} - \delta\frac{1}{h^2} + 2\alpha\frac{h}{y} + 3\gamma h^2\right)(h_\omega\omega_1 + h_w w_1),$$

or

$$\begin{aligned} h_{\omega yy} &= \left(2\frac{H}{h} - \frac{1}{y}\right) h_{\omega y} + \left(-\frac{H^2}{h^2} - \delta\frac{1}{h^2} + 2\alpha\frac{h}{y} + 3\gamma h^2\right) h_\omega, \\ h_{wyy} &= \left(2\frac{H}{h} - \frac{1}{y}\right) h_{wy} + \left(-\frac{H^2}{h^2} - \delta\frac{1}{h^2} + 2\alpha\frac{h}{y} + 3\gamma h^2\right) h_w. \end{aligned}$$

Hence the functions h_ω and h_w are a particular solutions of (39). It is easy to show that they are linearly independent, and so the general solution (39) has the form

$$F = h_\omega A(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n) + h_w B(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n). \tag{40}$$

Finally, from equations (37), (38), (40) it follows that the local higher $x-$symmetry Lie-Bäcklund of the system (34) is defined from

$$f = D^{-1}(h_\omega A + h_w B), \quad g = D^{-1}(h_{\omega y} A + h_{wy} B),$$

where functions $A = A(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n)$ and $B = B(x, \omega, w, \omega_1, w_1, \ldots, \omega_n, w_n)$ satisfy

$$\frac{\delta}{\delta\omega}(h_\omega A + h_w B) = \frac{\delta}{\delta w}(h_\omega A + h_w B) = 0.$$

The $y-$characteristic Lie ring of equations (34) is defined by the vector fields

$$Y_1 = \frac{\partial}{\partial p_1}, \quad Y_2 = \frac{\partial}{\partial q_1},$$

$$Y_3 = \frac{\partial}{\partial x} + p_1\frac{\partial}{\partial p} + q_1\frac{\partial}{\partial q} + q_1\frac{\partial}{\partial \bar{p}_1} + \left(\frac{q_1^2}{p_1} - \frac{q_1}{y} + \delta\frac{1}{p_1} + \beta\frac{1}{y} + \alpha\frac{p_1^2}{y} + \gamma p_1^3\right)\frac{\partial}{\partial \bar{q}_1} + \ldots.$$

It is easy to see that $y-$ring is infinite. However, the system of equations (34) has $y-$integral of the first order $\bar{\omega} = \bar{p}_1 - q$. Define higher $y-$symmetries

$$\varphi = \varphi(x, y, p, q, \bar{p}_1, \bar{q}_1, \ldots, \bar{p}_n, \bar{q}_n), \quad \psi = \psi(x, y, p, q, \bar{p}_1, \bar{q}_1, \ldots, \bar{p}_m, \bar{q}_m),$$
$$(p_\tau = \varphi,\ q_\tau = \psi)$$

for equations (34).

Let the order of variables $p, q, \bar{p}_1, \bar{q}_1, \bar{p}_2, \bar{q}_2, \ldots$ the functions $D\varphi$ and $D\psi$ be equal to n and m correspondingly. Then, from the defining system of equations

$$D\bar{D}\varphi = D\psi, \quad D\bar{D}\psi = \left(2\frac{q_1}{p_1} - \frac{1}{y}\right) D\psi + \left(-\frac{q_1^2}{p_1^2} - \delta\frac{1}{p_1^2} + 2\alpha\frac{p_1}{y} + 3\gamma p_1^2\right) D\varphi \tag{41}$$

it follows that $n+1 = m$ and $m+1 = m$, hence

$$D\varphi = F(x, y, p_1, q_1), \quad D\psi = G(x, y, p_1, q_1).$$

On the other hand

$$D\varphi(x, y, p, q, \bar{p}_1, \bar{q}_1, \ldots, \bar{p}_n, \bar{q}_n) = \frac{\partial\varphi}{\partial x} + p_1\frac{\partial\varphi}{\partial p} + q_1\left(\frac{\partial}{\partial\bar{p}_1} + \frac{\partial}{\partial q}\right)\varphi +$$

$$+ \left(\frac{q_1^2}{p_1} - \frac{q_1}{y} + \delta\frac{1}{p_1} + \beta\frac{1}{y} + \alpha\frac{p_1^2}{y} + \gamma p_1^3\right)\left(\frac{\partial}{\partial\bar{p}_2} + \frac{\partial}{\partial\bar{q}_1}\right)\varphi + \ldots.$$

Therefore, the function φ satisfies the system

$$\frac{\partial\varphi}{\partial x} = \alpha(x, y), \ \frac{\partial\varphi}{\partial p} = \alpha_0(x, y), \ \left(\frac{\partial}{\partial\bar{p}_1} + \frac{\partial}{\partial q}\right)\varphi = \alpha_1(x, y),$$

$$\left(\frac{\partial}{\partial\bar{p}_2} + \frac{\partial}{\partial\bar{q}_1}\right)\varphi = \alpha_2(x, y), \ldots, \ \left(\frac{\partial}{\partial\bar{p}_n} + \frac{\partial}{\partial\bar{q}_{n-1}}\right)\varphi = \alpha_n(x, y).$$

It is easy to obtain from it, that

$$\varphi = \beta(x, y) + \beta_0(y)p + \beta_1(y)\bar{p}_1 + \ldots + \beta_n(y)\bar{p}_n + h(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{n-1}). \tag{42}$$

An analogous formula holds true for the function ψ :

$$\psi = \gamma(x, y) + \gamma_0(y)p + \gamma_1(y)\bar{p}_1 + \ldots + \gamma_m(y)\bar{p}_m + H(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{m-1}). \tag{43}$$

Since the functions $\varphi = h(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{n-1})$ and $\psi = H(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{m-1})$ are symmetries of the system (34) for any h and H, then we obtain from (42) and (43), that

$$\varphi = \beta(x, y) + \sum_{k=0}^{n}\beta_k(y)\bar{p}_k, \quad \psi = \gamma(x, y) + \sum_{k=0}^{m}\gamma_k(y)\bar{p}_k \tag{44}$$

are also symmetries. Substituting (44) into defining system (41), we find that

$$\varphi = \varphi(y), \quad \psi = \psi(y).$$

Thus, $y-$symmetry of the system of equations (34) are calculate by formulae

$$\varphi = h(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{n-1}), \quad \psi = H(y, \bar{\omega}, \bar{\omega}_1, \ldots, \bar{\omega}_{m-1}).$$

Acknowledgments

This work is supported by the Russian Foundation for Basic Research (RFBR) grants 10-01-00088-a and 11-01-97005-r-povolzhye-a and by Federal Task Program "Scientic and pedagogical staff of innovative Russia for 2009-2013 contract no. 2012-1.5-12-000-1003-011.

References

[1] Kostrigina O.S., Zhiber A.V., *Darboux-integrable two-component nonlinear hyperbolic systems of equations*, J. Math. Phys. **52**, 033503 (2011); doi:10.1063/1.3559134 (32 pages).

[2] Goursat E., *Recherches sur quelques équations aux dérivées partielles du second ordre*, Annales de la faculté des Sciences de I'Université de Toulouse 2^e série, tome 1, n^0 1 (1899) p.31-78

[3] Zhiber A.V., Murtazina R.D., *On the characteristic Lie algebras for the equations* $u_{xy} = f(u, u_x)$, Fundam. Prikl. Mat. **12** (2006) 65-78 (in Russian)

[4] Zhiber A.V., Murtazina R.D., Habibullin I.T., Shabat A.B., *Characteristic Lie rings and integrable models in mathematical physics*, Ufimskii Matem. Zhurnal, v.4, № 3 (2012), pp. 17-85.

[5] Gürser M., Zhiber A.V., Habibullin I.T., *Characteristic Lie rings of differential equations*, Ufimskii Matem. Zhurnal, v.4, № 1 (2012), pp. 53-62.